

On a Nonlinear System of Partial Differential Equations

NIRZI G. DE ANDRADE

Instituto de Matemática, UFRJ, Rio de Janeiro, Brasil

Submitted by J. L. Lions

We present in this paper some results concerning the following nonlinear system of P.D.E.

$$(P) \begin{cases} u'' - cv'' + \alpha D^4 u - M(a(u)) D^2 u = f_1(x, t) & (1) \\ -cu'' + \gamma v'' + \delta D^4 v - \beta_0 D^2 v = f_2(x, t), & (2) \end{cases}$$

where $z' = z_t$, $Dz = z_x$ and $a(u) = \int_0^1 |Du|^2 dx$.

The above system is a mathematical model which describes coupled flexural and torsional oscillations of an open cross-section beam. In Part I we consider the abstract initial value problem associated with the above system, prove the existence and uniqueness of solutions in a weak sense and mention two applications. In Part II we obtain regular solutions when adequate conditions on the data are assumed.

INTRODUCTION

A particular case of system (P), namely, when $M(\lambda) = m_0$ ($m_0 > 0$) appears in Timoshenko *et al.* [13] as a mathematical model to describe coupled flexural and torsional vibrations of an open cross-section beam under certain conditions on longitudinal symmetry of the beam and on the load distribution.

The nonlinear term, which contains $\lambda = \int_0^L |Du|^2 dx$, appears when the extensibility of the beam is taken into account (see, for example, Ball [1]). Dickey [3] assuming existence of solutions, in Fourier series form, solved a mixed problem associated with Eq. (1) alone, with homogeneous boundary conditions, in the case $M(\lambda) = m_0 + m_1 \lambda$, $m_0 > 0$, $c = \alpha = 0$, $f_1 = 0$. Later, Dickey [4] considered the same equation but with $\alpha \neq 0$. The results of Dickey were generalized by Menzala [9] and Rivera [11]. A similar equation was also studied by Pohojaev [10]. Lions [6] studied the abstract problem associated with Eq. (1) in the case $c = \alpha = 0$ and $M = M(\lambda)$ of class $C^1(\mathbb{R}^+)$ and strictly positive. Medeiros [7] solved the same problem but

admitted the nonlinear term of the form $(\alpha + M(\lambda)) Du$ with $M \in C^1(\mathbb{R}^+)$, $M(\lambda) \geq m_0 + m_1 \lambda$, $\lambda \geq 0$, $\alpha \in \mathbb{R}$ and obtained solutions in a Gevrey class.

Ball in [1] obtained weak and classical solutions of Eq. (1) in the case $c = 0$ and $M(\lambda) = m_0 + m_1 \lambda$, $m_0 > 0$. Brito [2] studied an abstract problem related to Eq. (1) when $c = 0$ and rotary inertia is taken into account.

We shall consider $M(\lambda) = m_0 + m_1 \lambda$, $m_1 > 0$. The results, however, are still valid if $M = M(\lambda) \in C^1(\mathbb{R}^+)$, $M(\lambda) \geq m_0 > 0$.

The coefficients in Eqs. (1) and (2) are constants and represent quantities which have to do with the physical model. In particular, the constant c represents the distance from the centroidal to the shear center axis of the beam (see Timoshenko *et al.* [13]). The two constants γ and c are related by $\gamma = I_p/S + c^2$, where S and I_p denotes the cross-section area and the polar moment of inertia, respectively, and so we have $\gamma > c^2$.

THE ABSTRACT INITIAL VALUE PROBLEM

We consider, in this section, the abstract initial value problem associated with system P.

Let H be a Hilbert space with scalar product and norm denoted respectively by (\cdot, \cdot) and $|\cdot|$. Let A be a linear map in H from domain $D(A) \subset H$ into H . We shall also suppose that A is selfadjoint and positive. With the graph norm $D(A)$ is a Hilbert space which we denote by V . It follows that V is continuously embedded in H . We shall also admit that the immersion of V in H is compact and that V is separable. We define A^2 by

$$\langle A^2 u, v \rangle = (Au, Av),$$

where V' is the dual of V and $\langle \cdot, \cdot \rangle$ denotes the duality pairing $V'V$. It follows that $A^2 \in \mathcal{L}(V, V')$. If $u, v \in V$ let $a(u, v)$ and $b(u, v)$ be the bilinear forms given by

$$a(u, v) = (Au, v) \quad \text{and} \quad b(u, v) = (Au, Av).$$

If $A: D(A) \subset H \rightarrow H$ satisfies the above conditions and $u_0, v_0, u_1, v_1, f_1, f_2$ are given functions we consider the problem of finding functions u and v solutions of the system

$$u'' + cv'' + \alpha A^2 u + (m_0 + m_1 |A^{1/2} u|^2) Au = f_1 \quad (3)$$

$$cu'' + \gamma v'' + \delta A^2 v + \beta_0 Av = f_2 \quad (4)$$

verifying the initial conditions

$$u(0) = u_0, \quad v(0) = v_0; \quad u'(0) = u_1, \quad v'(0) = v_1. \quad (5)$$

THEOREM 1. Let $u_0, v_0 \in V, u_1, v_1 \in H, f_1, f_2 \in L^2(0, T; H)$ be given. Then a unique pair of functions $\{u, v\}$ exists defined in $[0, T]$ ($0 < T < +\infty$) with values in H such that:

$$(1) \quad u, v \in L^\infty(0, T; V),$$

$$(2) \quad u', v' \in L^\infty(0, T; H).$$

The functions u, v are solutions of the system

$$\begin{aligned} & \frac{d}{dt}(u'(t), \omega) + c \frac{d}{dt}(v'(t), \omega) + \alpha(Au(t), A\omega) \\ & + (m_0 + m_1 a(u))(Au(t), \omega) = (f_1(t), \omega), \\ (3) \quad & c \frac{d}{dt}(u'(t), \omega) + \gamma \frac{d}{dt}(v'(t), \omega) + \delta(Av(t), A\omega) \\ & + \beta_0(Av(t), \omega) = (f_2(t), \omega), \end{aligned}$$

in the sense of $\mathcal{D}'(0, T), \forall w \in V$.

$$(4) \quad u(0) = u_0, v(0) = v_0; u'(0) = u_1, v'(0) = v_1.$$

Proof of Theorem 1 (a) Existence of solutions. Since we suppose V separable there are vectors w_1, w_2, \dots, w_k in H such that the set $\{w_1, w_2, \dots, w_k\}$ is linearly independent for each $k \in N$ and the set of finite linear combinations of w_i are dense in V . We denote by V_m the subspace of V spanned by the first m vector w_1, w_2, \dots, w_m . We shall look for approximate solutions of system (3) of the form

$$u_m(t) = \sum_{j=1}^m g_{mj}(t) w_j; \quad v_m(t) = \sum_{j=1}^m h_{mj}(t) w_j,$$

that is, $u_m, v_m \in V_m$.

The two functions u_m, v_m shall be determined so that they are solutions of the approximate system:

$$\begin{aligned} & (u_m''(t), w) + c(v_m''(t), w) + \alpha(Au_m(t), Aw) + (m_0 + m_1 a(u_m))(Au_m(t), w) \\ & = (f_1(t), w), \end{aligned} \tag{6}$$

$$\begin{aligned} & c(u_m''(t), w) + \gamma(v_m''(t), w) + \delta(Av_m(t), Aw) + \beta_0(Av_m(t), w) \\ & = (f_2(t), w), \end{aligned} \tag{7}$$

$\forall w \in V_m$ and satisfy the initial conditions

$$u_m(0) = u_{0m}, \quad v_m(0) = v_{0m}; \quad u_m'(0) = u_{1m}, \quad v_m'(0) = v_{1m} \tag{8}$$

with

$$\begin{aligned} u_{0m} &\rightarrow u_0, & v_{0m} &\rightarrow v_0 & \text{strongly in } V, \\ u_{1m} &\rightarrow u_1, & v_{1m} &\rightarrow v_1 & \text{strongly in } H. \end{aligned}$$

The system (6–8) of ordinary differential equations has unique solution, for each $m \in N$, in an interval $[0, t_m)$. The a priori estimates that follow show, in particular, that $t_m = T$, $0 < T < +\infty$. The method used to get these a priori estimates is standard in this type of problem so we omit the details.

Taking $w = 2u'_m(t)$ in Eq. (6) and $w = 2v'_m(t)$ in Eq. (7) we obtain

$$\begin{aligned} &\frac{d}{dt} |u'_m|^2 + \alpha \frac{d}{dt} |Au_m|^2 + m_0 \frac{d}{dt} |A^{1/2}u_m|^2 + \frac{m_1}{2} \frac{d}{dt} |A^{1/2}u_m|^4 \\ &= 2(f_1, u'_m) - 2c(v''_m, u'_m), \\ &\gamma \frac{d}{dt} |v'_m|^2 + \delta \frac{d}{dt} |Av_m|^2 + \beta_0 \frac{d}{dt} |A^{1/2}v_m|^2 \\ &= 2(f_2, v'_m) - 2c(u''_m, v'_m). \end{aligned}$$

Adding the two equations and integrating in $(0, t)$, $t < t_m$, it follows that

$$\begin{aligned} &|u'_m|^2 + \gamma |v'_m|^2 + \alpha |Au_m|^2 + \delta |Av_m|^2 + m_0 |A^{1/2}u_m|^2 \\ &+ \beta_0 |A^{1/2}v_m|^2 + \frac{1}{2}m_1 |A^{1/2}u_m|^4 \\ &\leq C + 2 \int_0^t |(f_1, u'_m)| ds + 2 \int_0^t |(f_2, v'_m)| ds + 2c |(u'_m, v'_m)|, \end{aligned}$$

where C is a positive constant independent of m that bounds the terms involving the initial conditions. From now on C will denote a generic positive constant. Let $c_1 = \frac{1}{2}(1 - (c^2/\gamma))$ and $c_2 = \frac{1}{2}(\gamma - c^2)$. Then it follows from the last inequality that

$$\begin{aligned} &c_1 |u'_m|^2 + c_2 |v'_m|^2 + \alpha |Au_m|^2 + \delta |Av_m|^2 + m_0 |A^{1/2}u_m|^2 + \beta_0 |A^{1/2}v_m|^2 \\ &\leq C + C \int_0^t \{|u'_m|^2 + |v'_m|^2\} ds. \end{aligned}$$

By the Gronwall inequality we have

$$|u_m|, |v_m|, |u'_m|, |v'_m| \leq C, \quad |Au_m|, |Av_m| \leq C, \quad |A^{1/2}u_m|, |A^{1/2}v_m| \leq C, \quad (9)$$

where C is independent of m . So the solutions may, indeed, be extended to an interval $[0, T)$, $T < +\infty$. Moreover, the sequences (u_m) , (v_m) belong to

bounded sets in $L^\infty(0, T; V)$ and $(u'_m), (v'_m), (Au_m), (Av_m), (A^{1/2}u_m), (A^{1/2}v_m)$ are in bounded sets in $L^\infty(0, T; H)$. Taking subsequences $u_{r'}, v_{r'}$ it follows they converge weak-star to u and v , respectively, in $L^\infty(0, T; V)$ and $u'_{r'}, v'_{r'}, Au_{r'}, Av_{r'}, A^{1/2}u_{r'}, A^{1/2}v_{r'}$ converge weak-star to $u', v', Au, Av, A^{1/2}u$ and $A^{1/2}v$, respectively, in $L^\infty(0, T; H)$. By hypothesis the immersion of V in H is compact so it follows that $u_{r'}$ converges strongly to u in $L^2(0, T; H)$ (Lions [5]).

As to the nonlinear term let $J(u_m) = a(u_m)Au_m$. Using the estimates we obtained, it results that $(J(u_m))$ belongs to a bounded set in $L^\infty(0, T; H)$. So we can take a subsequence $(J(v_{r'}))$ that converges weak-star to a function ξ in $L^\infty(0, T; H)$. By monotonicity or by an argument used by Medeiros [8] or Ball [1] we obtain that $\xi = J(u) = a(u)Au$.

In (6) and (7) we write $u_{r'}$ and $v_{r'}$ instead of u_m and v_m and consider $r' > m$, m now a fixed natural number. We pass to the limit as $r' \rightarrow \infty$ and conclude that u, v satisfies condition (3) of the theorem. To verify the initial conditions we observe that $u, v \in L^\infty(0, T; v)$ and $u', v' \in L^\infty(0, T; H)$. Using the fact that u, v are solutions it can be shown that $u'', v'' \in L^\infty(0, T; V')$. So we can identify u and v with element of $C([0, T]; H)$ and u', v' with elements of $C([0, T]; V')$. Therefore $u(0), v(0), u'(0), v'(0)$ are meaningful.

(b) Uniqueness. Suppose there are two pairs of solutions $\{u, v\}$ and $\{\hat{u}, \hat{v}\}$ and set $u = \bar{u} - \hat{u}$, $v = \bar{v} - \hat{v}$. Then u, v satisfies the system

$$u'' + cv'' + \alpha A^2 u + m_0 Au + m_1 [a(\bar{u})A\bar{u} - a(\hat{u})A\hat{u}] = 0 \quad (10)$$

$$cu'' + \gamma v'' + \delta A^2 v + \beta_0 Av = 0 \quad (11)$$

and initial conditions

$$u(0) = v(0) = 0; \quad u'(0) = v'(0) = 0.$$

Following Lions [5] we define z and w by

$$z(t) = \int_t^s u(\sigma) d\sigma \quad \text{if } t \leq s \quad \text{and} \quad z(t) = 0 \quad \text{if } t > s,$$

$$w(t) = \int_t^s v(\sigma) d\sigma \quad \text{if } t \leq s \quad \text{and} \quad w(t) = 0 \quad \text{if } t > s,$$

and set

$$z_1(t) = \int_0^t u(\sigma) d\sigma; \quad w_1(t) = \int_0^t v(\sigma) d\sigma.$$

It follows that

$$\begin{aligned} z(t) &= z_1(s) - z_1(t), & w(t) &= w_1(s) - w_1(t), \\ z'(t) &= -u(t), & w'(t) &= -v(t), \\ z(s) &= 0, & w(s) &= 0, \\ z(0) &= z_1(s), & w(0) &= w_1(s). \end{aligned}$$

From (10) and (11) we have that

$$\begin{aligned} & \int_0^s \{ \langle u'', z \rangle + c \langle v'', z \rangle + \alpha(Au, Az) + m_0(Au, z) \\ & \quad + m_1(a(\bar{u})A\bar{u} - a(\hat{u})A\hat{u}, z) \} d\sigma = 0 \\ & \int_0^s \{ c \langle u'', w \rangle + \gamma \langle v'', w \rangle + \delta(Av, Aw) + \beta_0(Av, w) \} d\sigma = 0. \end{aligned}$$

From these two relations we obtain, after some algebraic calculations,

$$\begin{aligned} & |u(s)|^2 + \gamma |v(s)|^2 + \alpha |Az_1(s)|^2 + m_0 a(z_1(s)) + \beta_0 a(w_1(s)) \\ & \leq 2c |u(s)| |v(s)| + 2m_1 \int_0^s |(a(\bar{u})A\bar{u} - a(\hat{u})A\hat{u}, z(\sigma))| d\sigma. \end{aligned}$$

Using estimates (9) for \bar{u} and \hat{u} it follows that the integrand of the integral of the last inequality is bounded by $C|u|^2 + C|Az|^2$. Also, from the properties of $z = z(t)$ we have

$$|Az(\sigma)|^2 \leq 2\{|Az_1(s)|^2 + |Az_1(\sigma)|^2\},$$

and the last inequality becomes

$$\begin{aligned} & C_1 |u(s)|^2 + C_2 |v(s)|^2 + (\alpha - 2Cs) |Az_1(s)|^2 + m_0 a(z_1(s)) + \beta_0 a(w_1(s)) \\ & \leq 2C \int_0^s \{|u(\sigma)|^2 + |Az_1(\sigma)|^2\} d\sigma. \end{aligned}$$

Given ε_0 , $0 < \varepsilon_0 < \alpha$, let us choose s_0 such that $s_0 > 0$ and $\alpha - 2Cs_0 > \varepsilon_0$. Therefore, for all $s \in [0, s_0]$ we have $\alpha - 2Cs > \alpha - 2Cs_0 > \varepsilon_0$. With s_0 so chosen it follows that

$$|u(s)|^2 + |v(s)|^2 + \varepsilon_0 |Az_1(s)|^2 \leq C \int_0^s \{|u(\sigma)|^2 + |Az_1(\sigma)|^2\} d\sigma.$$

This implies, by the Gronwall inequality, that $u(s) = 0$ and $v(s) = 0$ for all $s \in [0, s_0]$. By a standard argument we conclude that $u = 0$, $v = 0$ in $[0, T]$.

Applications

Let $\Omega \subset \mathbb{R}^n$ be an open bounded set with regular boundary $\partial\Omega$, $A = -\Delta$ and $a(u) = |\nabla u|^2$.

(a) Let $H = L^2(\Omega)$, $V = H_0^2(\Omega)$, $V' = H^{-2}(\Omega)$. If $u_0, v_0 \in V$, $u_1, v_1 \in H$, $f_1, f_2 \in L^2(0, T; H)$ are given functions, then there is a unique pair of functions $u, v \in L^\infty(0, T; V)$, $u', v' \in L^\infty(0, T; H)$ and

$$\begin{aligned} u'' - cv'' + \alpha \Delta^2 u - \left(m_0 + m_1 \int_{\Omega} |\nabla u|^2 dx \right) \Delta u &= f_1 \\ cu'' + \gamma v'' + \delta \Delta^2 v - \beta_0 \Delta v &= f_2 \end{aligned} \quad (13)$$

$$u(0) = u_0, \quad v(0) = v_0; \quad u'(0) = u_1, \quad v'(0) = v_1$$

When $n = 1$, $\Omega = (0, L)$, $A = -\partial^2/\partial x^2$ this corresponds to the case of a beam with fixed end points which cannot rotate.

(b) Now let $H = L^2(\Omega)$, $V = H_0^1(\Omega) \cap H^2(\Omega)$. Hypothesis on A are satisfied. Let $u_0, v_0 \in V$, $u_1, v_1 \in H$, $f_1, f_2 \in L^2(0, T; H)$. Then there exists a unique pair of functions u, v solutions of problem (13).

In the case $n = 1$, $\Omega = (0, L)$, $A = -\partial^2/\partial x^2$ this corresponds to a beam in which the linear displacement and the bending moments are zero at the end points. (See Timoshenko [12]).

REGULAR SOLUTIONS

Let us consider again the system (3) – (4) with initial conditions (5) and suppose now that A is a linear operator in H , with dense domain $D(A)$, A selfadjoint and positive. We suppose also that there are sequences of numbers $\{\lambda_i\}_{i \in \mathbb{N}}$ and vectors $\{\omega_i\}_{i \in \mathbb{N}}$ such that $A\omega_j = \lambda_j\omega_j$, $j \in \mathbb{N}$, $0 < \lambda_1 < \lambda_2 < \dots < \lambda_n < \dots \rightarrow \infty$ and the set of eigenvectors is dense in H .

Now $u(t)$, $v(t)$ are vectors of H , t a real variable, and the derivative of $u(v)$ with respect to the norm of H denoted by u' (v') or du/dt (dv/dt). As before (\cdot, \cdot) and $|\cdot|$ denotes the scalar product and norm in H . We define $a(u, v) = (Au, v)$ and $a(u) = (Au, u)$, $u \in D(A)$.

Regular solutions are shown to exist according to the following theorem.

THEOREM 2. *Suppose that the operator A satisfies the above conditions and that $u_0, v_0, u_1, v_1, f_1, f_2$ are given functions such that*

- (1) $u_0, v_0 \in D(A^{k+1/2})$
- (2) $u_1, v_1 \in D(A^k)$
- (3) $f_1, f_2 \in D(A^k)$; $A^k f_1, A^k f_2 \in L^2(0, T; H)$ for all $k = 0, 1, 2, \dots$

Then there is a unique pair of functions u, v from $[0, T] \rightarrow H$, $0 < T < +\infty$ satisfying the conditions:

- (4) $u(t), v(t), u'(t), v'(t) \in D(A^k)$
- (5) $u, v, u', v' \in C^0([0, T]; D(A^k))$ for all $k = 0, 1, 2, \dots$
- (6) u, v satisfies system (3)–(4) and initial conditions (5).

Proof. (a) Existence of Regular Solutions. Let $V_m = [w_1, w_2, \dots, w_m]$ be the subspace of H generated by the m first eigenvectors of A and let u_m, v_m be defined by $u_m(t) = \sum_{i=1}^m g_{im}(t)w_i$, $v_m(t) = \sum_{i=1}^m h_{im}(t)w_i$. Functions $u_m, v_m \in V_m$ are to be solutions of the approximate system

$$(u_m''(t), w) + c(v_m''(t), w) + \alpha(Au_m(t), Aw) + (m_0 + m_1 a(u_m(t)))(Au_m(t), w) = (f_1, w) \quad (6')$$

$$c(u_m''(t), w) + \gamma(v_m''(t), w) + \delta(Av_m(t), Aw) + \beta_0(Av_m(t), w) = (f_2, w), \quad (7')$$

$\forall w \in V_m$ and satisfying the initial conditions

$$u_m(0) = u_{0m}, \quad v_m(0) = v_{0m}; \quad u_m'(0) = u_{1m}, \quad v_m'(0) = v_{1m}, \quad (8')$$

where

$$u_{\alpha m} = \sum_{i=1}^m (u_{\alpha}, w_i)w_i, \quad \alpha = 0, 1.$$

This system of ordinary differential equations has unique solutions u_m, v_m in an interval $(0, t_m)$, for each $m \in \mathbb{N}$.

Estimates 1

Proceeding exactly in the same way as we did in Part 1 we get the a priori estimates (9) for $u_m, v_m, u_m', v_m', Au_m, Av_m, A^{1/2}u_m$, and $A^{1/2}v_m$. These estimates permit us to extend u_m, v_m to the interval $[0, T]$, $T < +\infty$, T independent of m .

Estimates 2

Now let $w = 2A^{2k}u_m'$ in (6') and $w = 2A^{2k}v_m'$ in (7'). Taking into account the properties of A , the hypothesis on the initial data and on f_1, f_2 we obtain

$$\begin{aligned} & \frac{d}{dt} \{ |A^k u_m'|^2 + |A^{k+1} u_m|^2 + [m_0 + m_1 a(u_m)] |A^{k+1/2} u_m|^2 \} \\ &= 2(A^k f_1, A^k u_m') - 2c(A^k u_m'', A^k u_m') + 2m_1 |A^{k+1/2} u_m|^2 \frac{d}{dt} a(u_m). \end{aligned}$$

$$\begin{aligned} & \frac{d}{dt} \{ \gamma |A^k v_m'|^2 + \delta |A^{k+1} v_m|^2 + \beta_0 |A^{k+1/2} v_m|^2 \} \\ &= 2(A^k f_2, A^k v_m') - 2c(A^k u_m'', A^k v_m'). \end{aligned}$$

Adding the two equations, integrating and bounding the terms on the second member the following inequality is obtained:

$$\begin{aligned} & c_1 |A^k u'_m|^2 + c_2 |A^k v'_m|^2 + \alpha |A^{k+1} u_m|^2 + \delta |A^{k+1} v_m|^2 + m_0 |A^{k+1/2} u_m|^2 \\ & \quad + \beta_0 |A^{k+1/2} v_m|^2 \\ & \leq C \int_0^t \{ |A^k u'_m|^2 + |A^{k+1/2} u_m|^2 \} ds, \end{aligned} \quad (14)$$

where, as before, $c_1 = \frac{1}{2}(1 - (c^2/\gamma))$, $c_2 = \frac{1}{2}(\gamma - c^2)$.

From (14) and the Gronwall inequality it follows that

$$\begin{aligned} & |A^k u'_m|^2 + |A^k v'_m|^2 + |A^{k+1} u_m|^2 + |A^{k+1} v_m|^2 + |A^{k+1/2} u_m|^2 \\ & \quad + |A^{k+1/2} v_m|^2 \leq C \end{aligned} \quad (15)$$

in $(0, T)$, C independent of m .

Limit of Approximate Solutions

We shall prove first that the sequence $(A^k u'_m)$, $(A^k v'_m)$, $(A^{k+1/2} u_m)$, $(A^{k+1/2} v_m)$, $(A^{k+1} u_m)$, $(A^{k+1} v_m)$ are Cauchy sequences in $[0, T]$, $T < +\infty$. Suppose that u_{n_2} , u_{n_1} , v_{n_2} , v_{n_1} are approximate solutions of system (6'), (7') with initial conditions (8'). Let n_2 , n_1 be natural numbers and $n_2 > n_1$. Set $w = u_{n_2} - u_{n_1}$ and $z = v_{n_2} - v_{n_1}$. Then for all $\eta \in V_m$ ($m \geq n_2$) w , z satisfies the following two relations:

$$\begin{aligned} & (w'', \eta) + c(z'', \eta) + \alpha(Aw, A\eta) + m_0(Aw, \eta) + m_1 a(u_{n_1})(Aw, \eta) \\ & \quad + m_1 [a(u_{n_2}) - a(u_{n_1})] a(u_{n_2}, \eta) = 0 \end{aligned} \quad (16)$$

$$c(w'', \eta) + \gamma(z'', \eta) + \delta(Az, A\eta) + \beta_0(Az, \eta) = 0. \quad (17)$$

Taking $\eta = 2A^{2k} w'$ in (16) and $\eta = 2A^{2k} z'$ in (17). It follows that

$$\begin{aligned} & |A^k w'|^2 + \gamma |A^k z'|^2 + \alpha |A^{k+1} w|^2 + \delta |A^{k+1} z|^2 + m_0 |A^{k+1/2} w|^2 \\ & \quad + \beta_0 |A^{k+1/2} z|^2 \\ & \leq |K_k(0)| + 2c |A^k w'| |A^k z'| + 2m_1 C \int_0^t \{ |A^{k+1/2} w|^2 + |A^{k+1/2} z|^2 \\ & \quad + |A^k w'|^2 \} ds, \end{aligned} \quad (18)$$

where

$$\begin{aligned} K_k(0) = & |A^k w'(0)|^2 + \gamma |A^k z'(0)|^2 + \alpha |A^{k+1} w(0)|^2 + \delta |A^{k+1} z(0)|^2 \\ & + (m_0 + m_1 a(u_{n_2}(0))) |A^{k+1/2} w(0)|^2 + \beta_0 |A^{k+1/2} z(0)|^2 \\ & + 2c(A^k w'(0), A^k z'(0)). \end{aligned}$$

The hypothesis on the initial data imply that $K_k(0) \rightarrow 0$, as $n_2, n_1 \rightarrow \infty$, for all $k = 0, 1, 2, \dots$.

For $k = 0$ in (18) it follows that

$$\begin{aligned} & |w'|^2 + |z'|^2 + |Aw|^2 + |Az|^2 + |A^{1/2}w|^2 + |A^{1/2}z|^2 \\ & \leq C_0 K_0(0) + C \int_0^t \{|A^{1/2}w|^2 + |w'|^2\} ds \end{aligned}$$

hence by the Gronwall inequality

$$|A^{1/2}w|^2 \leq C_0 K_0(0) e^{CT}, \quad T < +\infty. \quad (19)$$

Taking (19) into (18) it follows that

$$\begin{aligned} & |A^k w'|^2 + |A^k z'|^2 + |A^{k+1}w|^2 + |A^{k+1}z|^2 + |A^{k+1/2}w|^2 + |A^{k+1/2}z|^2 \\ & \leq I_{0k}(T) + C \int_0^t \{|A^k w'|^2 + |A^{k+1/2}w|^2\} ds, \end{aligned}$$

where

$$I_{0k}(T) = C_0 K_k(0) + C C_0 K_0(0) T e^{CT}, \quad T < +\infty.$$

The last inequality implies that

$$\begin{aligned} & |A^k w'|^2 + |A^k z'|^2 + |A^{k+1}w|^2 + |A^{k+1}z|^2 + |A^{k+1/2}w|^2 + |A^{k+1/2}z|^2 \\ & \leq I_{0k}(T) e^{CT}, \quad \forall t \in [0, T]. \end{aligned}$$

But $\lim_{n_2, n_1 \rightarrow \infty} I_{0k}(T) = 0$, $\forall k \in \mathbb{N}$, so the sequences $(A^k u'_n(t))$, $(A^k v'_n(t))$, $(A^{k+1} u_n(t))$, $(A^{k+1} v_n(t))$, $(A^{k+1/2} u_n(t))$, $(A^{k+1/2} v_n(t))$ are uniform Cauchy sequences of $C^0([0, T], H)$. Therefore they are convergent for all $k \in \mathbb{N}$. Thus for $k = 0$ we have $u = u(t)$, $v = v(t)$ such that $(u_n(t))$, $(v_n(t))$ converge uniformly to $u(t)$, $v(t)$ in $C^0([0, T], H)$ and

$$u, v; u', v' \in C^0([0, T], D(A^k)) \quad \text{for all } k = 0, 1, 2, \dots$$

We also have the uniform convergence of $(A^{1/2} u_n)$ to $(A^{1/2} u)$ in $C^0([0, T], H)$ and hence we may pass to the limit in the nonlinear term. It follows from the approximate problem that

$$\begin{aligned} & \frac{d}{dt} (u'(t), w) + c \frac{d}{dt} (v'(t), w) + \alpha (Au(t), Aw) \\ & + (m_0 + m_1 a(u(t)))(Au(t), w) = (f_1, w) \\ & c \frac{d}{dt} (u'(t), w) + \gamma \frac{d}{dt} (v'(t), w) + \delta (Av(t), Aw) + \beta_0 (Av(t), w) \\ & = (f_2, w), \quad \forall w \in H. \end{aligned}$$

The uniform convergence of $(u_n(t))$, $(v_n(t))$, $(u'_n(t))$, $(v'_n(t))$ in $C^0([0, T]; H)$ guarantees that the initial conditions are satisfied.

Uniqueness

Suppose there are two pairs of regular solutions $\{u_1, v_1\}$, $\{u_2, v_2\}$ in $C^0([0, T]; H)$ of the system (3), (4) satisfying initial conditions (5). Let $w = u_2 - u_1$ and $z = v_2 - v_1$ and $v_2 - v_1$. Then $\{w, z\}$ satisfies the following two equations:

$$\begin{aligned} w'' + cz'' + \alpha A^2 w + m_0 A w + m_1 a(u_2) A w &= m_1 [a(u_1) - a(u_2)] A u_1 \\ cw'' + \gamma z'' + \delta A^2 z + \beta_0 A z &= 0 \end{aligned}$$

and the initial conditions

$$w(0) = z(0) = 0; \quad w'(0) = z'(0) = 0.$$

Taking the scalar product of the first equation with $2w'$ and of the second one with $2z'$, integrating in $[0, t]$, $t \leq T$, we get

$$\begin{aligned} &|w'|^2 + \gamma |z'|^2 + \alpha |Aw|^2 + |Az|^2 + m_0 |A^{1/2} w|^2 + \beta_0 |A^{1/2} z|^2 \\ &\leq 2c |w'| |z'| + 2m_1 \int_0^t |A^{1/2} w|^2 \left| \frac{d}{ds} a(u_2) \right| ds \\ &\quad + 2m_1 \int_0^t [a(u_2) - a(u_1)] |Au_1| |w'| ds. \end{aligned}$$

From this inequality and the estimates we obtained before, it follows that

$$\begin{aligned} &|w'|^2 + |z'|^2 + |Aw|^2 + |Az|^2 + |A^{1/2} w|^2 + |A^{1/2} z|^2 \\ &\leq C \int_0^t \{|w'|^2 + |Aw|^2 + |A^{1/2} w|^2\} ds \end{aligned}$$

and this implies that $w = 0$, $z = 0$ in $[0, T]$.

Applications

Let H be the Hilbert space $L^2(\Omega)$ for $\Omega \subset \mathbb{R}^n$ an open bounded set with regular boundary $\partial\Omega$. We take $A = -\Delta$, where Δ is the Laplacian operator, and consider the system:

$$\begin{aligned} u'' + cv'' + \alpha \Delta^2 u - \left(m_0 + m_1 \int_{\Omega} |\nabla u|^2 dx \right) \Delta u &= f_1 \\ cu'' + \gamma v'' + \delta \Delta^2 v - \beta_0 \Delta v &= f_2 \end{aligned} \quad (20)$$

together with the initial conditions

$$u(0) = u_0, \quad v(0) = v_0; \quad u'(0) = u_1, \quad v'(0) = v_1. \quad (21)$$

(a) We take $u_0, v_0 \in D(A^{k+1/2})$, $u_1, v_1 \in D(A^k)$ and $D(A) = H^4(\Omega) \cap H_0^2(\Omega)$.

Then $\{u, v\}$ is the unique solution of (2) satisfying initial conditions (21), $u, v \in C^0([0, T]; D(A^k))$.

(b) Let $u_0, v_0 \in D(A^{k+1/2})$, $u_1, v_1 \in D(A^k)$ and $D(A) = H^2(\Omega) \cap H_0^1(\Omega)$. Again, $\{u, v\}$ is the unique solution of (20) satisfying (21), $u, v \in C^0([0, T]; D(A^k))$. When $n = 1$ the interpretations are identical with those given in Part I.

We note again that these results remain valid if the nonlinear terms has for coefficient a function $M = M(\lambda)$, $\lambda \geq 0$, $M \in C^1$, and $M \geq m_0 > 0$.

ACKNOWLEDGMENT

This paper is part of my thesis written under the invaluable guidance of Prof. Luiz Adauto Medeiros who drew my attention to this problem and to whom I here express all my gratitude.

REFERENCES

1. J. M. BALL, Initial boundary value problems for an extensible beam, *J. Math. Anal. Appl.* **42** (1973), 61–90.
2. E. H. DE BRITO, A nonlinear hyperbolic equation, *Internat. J. Math. Math. Sci.* **3** (3) (1980), 505–520.
3. R. W. DICKEY, Infinite systems of nonlinear oscillations equations related to the string, *Proc. Amer. Math. Soc.* (1969), 459–468.
4. R. W. DICKEY, Free vibrations and dynamic buckling of the extensible beam, *J. Math. Anal. Appl.* **29** (1970), 443–454.
5. J. L. LIONS, "Quelques Méthodes de Résolution des Problèmes Aux Limites non Linéaires," Dunod, Paris, 1969.
6. J. L. LIONS, "On Some Questions in Boundary Value Problems of Mathematical Physics," Lecture Notes, Inst. Mat. Univ. Fed. do Rio de Janeiro, 1977.
7. L. A. MEDEIROS, On a new class of nonlinear wave equations, *J. Math. Anal. Appl.* **69** (1) (1979), 252–262.
8. L. A. MEDEIROS, "Weak Solutions for a Nonlinear Evolution Equation," to appear.
9. G. P. MENZALA, Une solution d'une équation nonlinéaire d'évolution, *C. R. Acad. Sci. Paris, Ser. A* **6**, 286.
10. S. I. POHOJAEV, On a class of quasilinear hyperbolic equations, *Math. USSR-Sb.* **25** (1), 145–158.
11. P. H. RIVERA, On a nonlinear hyperbolic equation in Hilbert spaces, *An. Acad. Brasil Ciênc.* **50** (2) (1978).
12. S. TIMOSHENKO, "Strength of Materials," 3rd ed., Part II, Van Nostrans-Reinhold, 1956.
13. S. TIMOSHENKO, D. H. YOUNG, AND W. WEAVER, "Vibration Problems in Engineering," 4th ed., Wiley, New York, 1974.